

AMENDMENTS TO THE CLAIMS

1. (currently amended) A method for selecting a set of normalizing data points from n data sets, where n is at least 3, containing data points having values and identities, the method comprising:
 - receiving n data sets;
 - considering the data points to be distributed in an n -dimensional data-point space;
 - determining one or more order-preserving sequences of data points within the n -dimensional data-point space; ~~and,~~
 - selecting, as normalizing data points, data points from the one or more order-preserving sequences; and
 - storing the selected normalizing points in a computer memory as a basis for subsequent normalization of the n data sets.
2. (original) The method of claim 1 wherein the one or more order-preserving sequences of data points is a single, longest order-preserving sequence of data points.
3. (original) The method of claim 1 wherein the data points within n data sets are associated with weights and wherein the one or more order-preserving sequences of data points is an order-preserving sequence of data points with a greatest sum of weights.
4. (original) The method of claim 1 wherein the one or more order-preserving sequences of data points is a longest order-preserving sequence of data points having a shortest Euclidian distance accumulated along a path from an initial data point of the order-preserving sequence to a final data point of the order-preserving sequence.
5. (original) The method of claim 1 wherein the one or more order-preserving sequences of data points are order-preserving sequences of data points of lengths within a threshold value of the length of an order-preserving sequence of data points of maximum length.

6. (original) The method of claim 1 wherein the data points within n data sets are associated with weights and wherein the one or more order-preserving sequences of data points are order-preserving sequences of data points with sums of weights within a threshold value of the sum of weights of an order-preserving sequence of data points with a greatest sum of weights.

7. (original) The method of claim 1 wherein considering the data points to be distributed in an n -dimensional data-point space further includes, for each data point, considering the data point to have a value in each of n -dimensions, the value of a data-point in an i th dimension equal to the value of the data point in an i th data set, where $1 \leq i \leq n$.

8. (original) The method of claim 1 wherein determining an order-preserving sequence of data points within the n -dimensional data-point space further includes:

- for each currently considered dimension,
 - ordering the data points with respect to the currently considered dimension;
 - traversing the ordered data points in a first direction, determining a metric corresponding to a maximum subsequence for each data point in the first direction; and
 - traversing the ordered data points in a second direction, determining a metric corresponding to a maximum subsequence for each data point in the second direction;
 - summing the determined metrics for each data point in each dimension to produce a metric sum for each data point; and
 - selecting as belonging to the maximum order-preserving sequence of data points those data points having a greatest metric sum.

9. (original) The method of claim 8 wherein selecting, as normalizing data points, data points from the order-preserving sequence further includes selecting data points with a metric sum greater than a threshold value.

10. (original) The method of claim 8 wherein selecting, as normalizing data points, data points from the one or more order-preserving sequences further includes selecting data points of a single order-preserving sequence.

11. (original) The method of claim 8 wherein selecting, as normalizing data points, data points from the one or more order-preserving sequences further includes selecting data points that most evenly partition the data points into subsets of data points.

12. (original) Computer instructions stored in a computer readable medium that implement the method of claim 1.

13. Cancelled

14. (currently amended) A system for selecting a set of normalizing data points from n data sets, where n is at least 3, containing data points having values and identities, the system comprising:

a processor;

a memory;

and computer instructions that select the set of normalizing data points from n data sets by

receiving n data sets,

considering the data points to be distributed in an n -dimensional data-point space,

determining one or more order-preserving sequence of data points within the n -dimensional data-point space, and

selecting, as normalizing data points, data points from the one or more order-preserving sequences; and

storing the selected normalizing points in a computer memory as a basis for subsequent normalization of the n data sets.

15. (currently amended) The ~~method~~ system of claim 14 wherein the one or more order-preserving sequences of data points is a single, longest order-preserving sequence of data points.

16. (currently amended) The ~~method~~ system of claim 14 wherein the data points within n data sets are associated with weights and wherein the one or more order-preserving sequences of data points is an order-preserving sequence of data points with a greatest sum of weights.

17. (currently amended) The ~~method~~ system of claim 14 wherein the one or more order-preserving sequences of data points is a longest order-preserving sequence of data points having a shortest Euclidian distance accumulated along a path from an initial data point of the order-preserving sequence to a final data point of the order-preserving sequence.

18. (currently amended) The ~~method~~ system of claim 14 wherein the one or more order-preserving sequences of data points are order-preserving sequence of data points within a threshold value of an order-preserving sequences of data points of maximum length.

19. (currently amended) The ~~method~~ system of claim 14 wherein the one or more order-preserving sequences of data points are order-preserving sequence of data points within a threshold value of an order-preserving sequences of data points with a greatest sum of weights.

20. (currently amended) A method for selecting a set of normalizing data points from n data sets, where n is at least 4 and even, containing data points having values and identities, the method comprising:

receiving n data sets;

considering the data points to be distributed in $\frac{n}{2}$ 2-dimensional data-point

spaces;

determining one or more order-preserving sequences of data points for each of the $\frac{n}{2}$ 2-dimensional data-point spaces; ~~and,~~
 selecting, as normalizing data points, data points from the order-preserving sequences; and
storing the selected normalizing points in a computer memory as a basis for subsequent normalization of the n data sets.

21. (original) The method of claim 20 wherein the one or more order-preserving sequences of data points is a single, longest order-preserving sequence of data points.

22. (original) The method of claim 20 wherein the data points within n data sets are associated with weights and wherein the one or more order-preserving sequences of data points is an order-preserving sequence of data points with a greatest sum of weights.

23. (original) The method of claim 20 wherein the one or more order-preserving sequences of data points is a longest order-preserving sequence of data points having a shortest Euclidian distance accumulated along a path from an initial data point of the order-preserving sequence to a final data point of the order-preserving sequence.

24. (original) The method of claim 20 wherein the one or more order-preserving sequences of data points are order-preserving sequences of data points within a threshold value of an order-preserving sequence of data points of maximum length.

25. (original) The method of claim 20 wherein the data points within n data sets are associated with weights and wherein the one or more order-preserving sequences of data points are order-preserving sequences of data points with sums of weights within a threshold value of the sum of weights of an order-preserving sequence of data points with a greatest sum of weights.

26. (original) The method of claim 20 wherein determining an order-preserving sequence of data points within a 2-dimensional data-point space further includes:

- for each currently considered dimension,
 - ordering the data points with respect to the currently considered dimension;
 - traversing the ordered data points in a first direction, determining a metric corresponding to a maximum subsequence for each data point in the first direction; and
 - traversing the ordered data points in a second direction, determining a metric corresponding to a maximum subsequence for each data point in the second direction;
 - summing the determined metrics for each data point in each dimension to produce a metric sum for each data point; and
 - selecting as belonging to the maximum order-preserving sequence of data points those data points having a greatest metric sum.

27. (original) The method of claim 20 wherein selecting, as normalizing data points, data points from the one or more order-preserving sequences further includes selecting data points which occur in the one or order-preserving sequences computed for greater than a threshold fraction of the $\frac{n}{2}$ 2-dimensional data-point spaces.

28. (original) Computer instructions stored in a computer readable medium that implement the method of claim 20.

29. Cancelled